



The future of zoological taxonomy is integrative, not minimalist

Alireza Zamani, Davide Dal Pos, Zdenek Faltýnek Fric, Alexander B. Orfinger, Mark D. Scherz, Alena Sucháčková Bartoňová & Hugo F. Gante

To cite this article: Alireza Zamani, Davide Dal Pos, Zdenek Faltýnek Fric, Alexander B. Orfinger, Mark D. Scherz, Alena Sucháčková Bartoňová & Hugo F. Gante (2022) The future of zoological taxonomy is integrative, not minimalist, *Systematics and Biodiversity*, 20:1, 1-14, DOI: [10.1080/14772000.2022.2063964](https://doi.org/10.1080/14772000.2022.2063964)

To link to this article: <https://doi.org/10.1080/14772000.2022.2063964>



Published online: 19 May 2022.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

PERSPECTIVE



The future of zoological taxonomy is integrative, not minimalist

ALIREZA ZAMANI¹ , DAVIDE DAL POS² , ZDENEK FALTÝNEK FRIC³ ,
ALEXANDER B. ORFINGER^{4,5} , MARK D. SCHERZ⁶ , ALENA SUCHÁČKOVÁ BARTOŇOVÁ³ 
& HUGO F. GANTE^{7,8,9} 

¹Zoological Museum, Biodiversity Unit, University of Turku, 20500 Turku, Finland

²Department of Biology, University of Central Florida, 4110 Libra dr. Rm 442, Orlando, FL 32816, USA

³Department of Biodiversity and Conservation Biology, Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Branisovska 31, Ceske Budejovice, CZ-37005, Czech Republic

⁴Department of Entomology and Nematology, University of Florida, Gainesville, FL 32611, USA

⁵Center for Water Resources, Florida A&M University, Tallahassee, FL 32301, USA

⁶Natural History Museum of Denmark, University of Copenhagen, Copenhagen, 1350, Denmark

⁷cE3c—Center for Ecology, Evolution and Environmental Changes, Universidade de Lisboa, Lisboa, Portugal

⁸Department of Biology, KU Leuven, Section Ecology, Evolution and Biodiversity Conservation, Charles Deberiotstraat 32 box 2439, Leuven, B-3000, Belgium

⁹Royal Museum for Central Africa, Leuvensesteenweg 17, Tervuren, 3080, Belgium

Roughly 18,000 species are described annually as new to science, while estimated extinction rates are comparable to or even exceeding these new discoveries. Considering the estimates of up to 15 million extant eukaryotic species on Earth, of which only about 2 million have been described so far, there has been a recent ‘boom’ of new potential approaches to more quickly discover and describe the millions of unknown species. This deficit is particularly noted in hyperdiverse taxa, as the current rate of species discovery is considered too slow. Recently, a ‘minimalist’ alpha taxonomic approach was proposed, relying solely on DNA barcoding and a habitus photograph, in a claimed effort to expedite the naming of new species to combat the so-called taxonomic impediment. In this paper, we point to limitations of minimalist taxonomy, present arguments in favour of the integrative approach, and finally explore a number of potential solutions to combat the taxonomic impediment in hyperdiverse taxa without sacrificing utility and quality for apparent speed and quantity.

Key words: automated morphological classification, biodiversity, citation mandate, eDNA, hyperdiverse taxa, integrative taxonomy, museomics, species description, systematics, taxonomic impediment

Introduction

It is estimated that there are up to 15 million extant eukaryotic species on Earth, of which only about 2 million have been discovered and described so far (Mora et al., 2011; Zamani et al., 2021). This puts taxonomists in a race against time to discover and describe the vast unknown portion of Earth’s biodiversity before it is lost as a result of the Anthropocene mass extinction. Each year about 18,000 species are described as new to science, with the annual species loss similar to, or even higher than, the rate of new species discoveries (Zamani et al., 2021), excluding the annual number of new synonyms which could be quite significant (e.g., around 20%

of names in insects; Stork, 1997). While naming new species is important, it is but one of the goals taxonomists are tasked with, the others being to circumscribe, properly describe and classify new taxa alongside their relatives (Seberg et al., 2003), as well as to curate existing names. Whereas naming follows enforced convention, species delimitation and their systematic relationships are scientific hypotheses rather than facts (Pante et al., 2015). As such, these practices need to be formulated in a way that clarifies species’ differences from other previously described species via a clear *diagnosis* (i.e., unique traits or combinations of traits), as well as a suite of shared traits representing the ancestor-descendant lines that connect all life (ICZN, 1999; Pante et al., 2015). This process can be time consuming, and new approaches are needed to quickly discover and

Correspondence to: Hugo F. Gante. E-mail: hugo.gante@kuleuven.be

accurately describe millions of currently unknown species in a timelier manner.

Since the foundational paper by Hebert *et al.* (2003) introducing a standardised fragment of mitochondrial cytochrome oxidase I (COI; *i.e.*, a molecular ‘barcode’) as a means of *identifying* known animal species, DNA barcoding has served a major role in biological sciences. In particular, DNA markers, especially DNA barcodes, have been a boon to species *discovery* when complemented with morphological studies and additional molecular and ecological data, made possible by having DNA barcodes of previously described species (*e.g.*, Vieites *et al.*, 2009). Its utility in systematics is widely recognised, and since its inception, DNA barcoding has been seen as a useful, albeit imperfect, tool in integrative taxonomy (*e.g.*, Dayrat, 2005; DeSalle *et al.*, 2005; Wheeler *et al.*, 2004). A recent move towards a ‘minimalist’ alpha taxonomic approach relies solely on a consensus DNA barcode and a habitus photograph (Meierotto *et al.*, 2019; Sharkey *et al.*, 2021b). This effort to expedite the *naming* of new wasp species to combat the so-called taxonomic impediment represents a problematic inflexion in the applications of DNA barcoding, contrary to principles professed by its own proponents when this discussion first emerged in the early 2000s. In several DNA barcoding proponents’ own words, “the discovery of previously undescribed species is augmented when information on morphological and ecological characters are fused with DNA barcodes” (Hebert & Barrett, 2005). Also, “we emphasize that DNA barcoding seeks merely to aid in *delimiting* species – to highlight genetically distinct groups exhibiting levels of sequence divergence suggestive of species status. By contrast, DNA barcodes – by themselves – are never sufficient to *describe* new species” (Hebert & Gregory, 2005). Finally, “it is important to note that it [DNA barcoding] does not seek to replace the Linnaean system of classification, and thus differs fundamentally from proposals to create a new taxonomic system based solely on DNA” (Gregory, 2005). “Hence, we urge strong collaborations between barcoders and taxonomists” (Hebert & Barrett, 2005).

This recent shift towards a DNA barcoding-based minimalist taxonomy materialises early concerns that it would try to replace an integrative taxonomy and classification system, as expressed by dozens of taxonomists and systematists (*e.g.*, Carvalho *et al.*, 2007; Ebach & Holdrege, 2005; Lipscomb *et al.*, 2003; Moritz & Cicero, 2004; Seberg *et al.*, 2003; Will *et al.*, 2005) and has received substantial criticism recently (Ahrens *et al.*, 2021; Engel *et al.*, 2021; Meier *et al.*, 2022; Zamani *et al.*, 2021, 2022). While we note the inconsistency above for historical context, our goal here is to highlight

why DNA taxonomy, and DNA barcoding in particular, is fraught with failure in resolving the taxonomic impediment of any group, particularly of hyperdiverse taxa. In turn, we present arguments in favour of the more robust integrative approach to taxonomy and systematics, and further explore potential solutions to the taxonomic impediment in hyperdiverse taxa, without sacrificing utility and quality of taxonomic descriptions for perceived speed and quantity.

Why (not) move toward a minimalist barcoding-based taxonomy?

Minimalist taxonomy’s reliance on a single mitochondrial barcode is at the crux of the problem. It has been known for decades that mitochondrial phylogenies frequently depart from the species tree for several reasons, such as introgression, incomplete lineage sorting, and infection with reproduction-manipulating endosymbionts in the case of arthropods and filarial nematodes (Funk & Omland, 2003). Even if consistent with the evolutionary history of a species, a mitochondrial barcode might be uninformative for delineating species, particularly in rapidly speciating, hyperdiverse groups (Wiemers & Fiedler, 2007). Thus, it comes without surprise that barcoding approaches that incorporate the multispecies coalescent into the analysis of multilocus sequence data have been proposed (Dowton *et al.*, 2014), even if these more robust statistical methods tend to over-split species in widespread taxa, especially if geographical sampling is inadequate (*e.g.*, Chambers & Hillis, 2020; Mason *et al.*, 2020). Because mitochondrial DNA has clear and long-known evolutionary blind spots, DNA barcoding as the sole source of information for species *delineation* and *description* does not serve taxonomy and should be discouraged.

The main argument put forward in favour of the minimalist approach is the unsubstantiated claim that it accelerates the naming of new species as a strategy to combat the taxonomic impediment (Sharkey *et al.*, 2021a). Nevertheless, formal analyses of the impacts of integrative taxonomy did not find a slowing effect of integrating across multiple types of data for species delimitation and description (Pante *et al.*, 2015). If anything, taxonomists should address bottlenecks imposed at different stages of the current taxonomic process, not replace it with subpar solutions. It is important to optimise taxonomic practice so that new descriptions are less likely to require revision in the foreseeable future, a goal that minimalist taxonomy is unwilling to achieve. Of note is that the proponents of the minimalist approach perform the minimum effort needed to barely conform to taxonomic naming conventions, leaving the

more time and effort consuming tasks of providing the integrative redescriptions to “future revisers” (Sharkey et al., 2021a). For many taxonomists, professionals and non-professionals alike, the ‘joy of discovery’ and naming of species are major driving forces keeping them active in such an underfunded field. Not only is taking away this small reward from these researchers unfair, in the long run it further harms taxonomy by discouraging taxonomists from conducting research on certain groups.

Is morphology dead?

The gold standard in taxonomic species descriptions is still morphology, despite a sustained increase over the years in integrative studies that also include other data types (Pante et al., 2015; Bond et al., 2021). Morphology helps us connect all life, former and current. Most species that have evolved are now extinct and many of these were diverse, ecologically important, and distinct from their living relatives (Wiens, 2004). This means that if we do not take fossil lineages into consideration, we would be ignoring over 99% of the taxa that have ever lived (Novacek & Wheeler, 1992). Minimalist taxonomy effectively excludes palaeontology from our study of evolutionary history and species relationships. As argued several times by Wheeler (2008; 2009; 2018), taxonomy, although essential for environmental biology, is at heart an evolutionary science. Taxonomy seeks to recognize evolutionary history, to describe what makes species unique (not just what makes them identifiable) and what they share with other species, and to phylogenetically classify them. These goals are not achievable based on DNA alone, and certainly not with single (or few) molecular markers.

Furthermore, while morphology is one *type* of data, different phenotypic traits are the manifestation of hundreds or thousands of genes, whose expression is channelled during development and filtered by ecological constraints of each species over evolutionary time. Thus, morphology is a rich source of information for robust taxonomic inference, although not without its own limitations. One of the frequent criticisms against morphology is the existence of cryptic species, which might reflect low resolution of traditional characters rather than a real absence of morphological differences should higher resolution methods be employed.

Integrative taxonomy is dead, long live integrative taxonomy!

What can be done to avoid one type of data from misrepresenting species diversity? The solution lies in integrative approaches. Olave et al. (2014) concluded that

without integrating multiple lines of evidence, interpreting what DNA-based approaches actually delimit will remain ambiguous. Integrative approaches narrow the parameter space where species are undetected in groups differentially diverging along different data type axes (Edwards & Knowles, 2014), and over-splitting is avoided especially if delimitations are congruent across different methods (Carstens et al., 2013). Furthermore, the use of integrative data also provides coherence between species detection and description (Edwards & Knowles, 2014), which should help ease the taxonomic impediment. There are multiple recent examples in which integrative strategies were successful at delimiting species, even in the case of complex evolutionary histories, morphological crypsis and hyperdiverse groups (e.g., Abdala et al., 2021; Arthofer et al., 2013; Brasero et al., 2020; Chaplin et al., 2020; Costa & Katz, 2021; Feliciano et al., 2021; Gebiola et al., 2012; Lima et al., 2020; Newton et al., 2020; Papakostas et al., 2016; Puillandre et al., 2012; Wachter et al., 2015; Weston et al., 2020). Furthermore, reanalysis of COI barcode clusters (“BINs”) computed by BOLD Systems (Ratnasingham & Hebert 2007) already indicate minimalist species are unstable, which would require redescription with additional evidence (Meier et al., 2022).

How would real-world examples have been affected by the minimalist approach?

Our objections to the minimalist approach are best illustrated by real-world examples. We argue that the approach produces species descriptions that are not usable by the taxonomic community, would alienate non-professional taxonomists (who represent the majority in some groups) and parataxonomists, does not address the true bottleneck of species discovery as opposed to species description, and in the long run would slow down species descriptions instead of speeding them up.

Much of the earlier discussion surrounding the use of DNA barcoding has occurred on theoretical grounds (e.g., Moritz & Cicero, 2004): what would the performance of DNA barcoding be, given the behaviour of mitochondrial DNA or the levels of divergence observed between species? How would DNA taxonomy impact species discoveries? We are now at a stage where we can directly assess the potential use of DNA barcoding in driving taxonomy, in particular in hyperdiverse groups. We take as our real-world examples tropical parasitoid wasps, African Great Lakes’ cichlids, and European butterflies. We feel these represent good test cases. Hyperdiverse taxa such as parasitoid wasps were the target group for the minimalist approach. Cichlids

and butterflies are well-studied, with barcode data available, which allow *de facto* statements about applicability of a DNA barcoding-based minimalist approach.

Ichneumon wasps. Consider Ichneumonidae as a group that contains more species than its sister family Braconidae. Dal Pos and Rouse (2018) described a new species of *Genaemirum* Heinrich, 1936 based on the dichotomous key by Rouse *et al.* (2016), which in turn was based (with modification) on the extensive work by Heinrich (1967), who provided the first and, so far, only taxonomic treatment of all Afrotropical genera of the subfamily Ichneumoninae (over 400 genera and 4300 species worldwide). Johansson (2020) described five new species of Diplazontinae from Sweden after Klopstein's (2014) revision of the group. Di Giovanni *et al.* (2018) identified a new species of the genus *Misetus* Wesmael, 1845 thanks to the key to the genera of Phaeogenini by Selfa and Diller (1994). Di Giovanni *et al.* (2015) and Di Giovanni and Reshchikov (2016) were able to fill the gaps of distribution of Ichneumonidae in Italy and record several species new to the country using the morphological keys and diagnoses provided by different European authors (e.g., Tereshkin, 2009). Moreover, many other taxonomic revisions are expected to be extremely useful in the years to come (e.g., Pham *et al.*, 2020; Rouse *et al.*, 2013). Sorting through an estimated >100,000 specimens of Ichneumonidae wasps from Uganda, morphological analysis of 456 individuals of the subfamily Rhyssinae resulted in a review of all Afrotropical rhyssines and the description of two new species (Hopkins *et al.*, 2019). The sorting into morphospecies was possible due to a previous review by Rouse and van Noort (2014) of 30 previously described individuals and twelve known species.

Without morphology, an identification key would be replaced by the necessity to DNA barcode all specimens. The alternative of barcoding just some specimens would require splitting them into preliminary morphospecies to select which specimens to barcode. This rather defeats the stated purpose of minimalist descriptions, which is to save time and effort. Based on the examples mentioned above, morphological keys can be extremely functional for identifying even hyperdiverse taxa, and the idea that they do not work lacks factual support. For example, it is doubtful Varga (2020) would have described a new Kenyan rhyssine species, had there not been multiple morphological characters and an identification key available. We do recognize that morphological keys are not always convenient, accurate, or easy to use, and do sometimes require specialised terminology and some expertise to understand and

implement. However, this is a good argument for integrating molecular data and investing in taxonomy and taxonomic training to improve both knowledge of species in such taxa, and the quality and ease of use of data to make keys more reliable, not as an excuse to abandon integrative studies. One recent example on how keys can be made more efficient is given by Fernandez-Triana (2022).

The central argument of increased speed has not been properly demonstrated by the minimalist taxonomy either. One important time sink pre-dates the act of describing species, and is 'hidden' in the form of sampling, building expertise and initial sorting of diversity usually based on morphology (known and unknown), as also recently noted but then not quantified by Fernandez-Triana (2022). For instance, it took Hopkins *et al.* (2019) several years to separate the parasitoid wasps, about a year to sort the rhyssines into species and write the descriptions. A *very* generous estimate would be that three months would have been saved using a minimalist approach, out of a total of at least three years at the cost of slowing down future work by established taxonomists and preventing new taxonomists from arising.

African Great Lakes' cichlids. The most spectacular examples of adaptive radiation in extant vertebrates are arguably the cichlids inhabiting the African Great Lakes (e.g., Gante & Salzburger, 2012). While the cichlid faunas from Lakes Malawi and Victoria are still mostly undescribed and number in the hundreds, the cichlid assemblage from Lake Tanganyika is comparatively well-known. In this older lake, the circa 250 species belong to 16 tribes defined on morphological grounds and confirmed by molecular data. These species radiated *in situ* from a common ancestor about 9.7 Ma (Ronco *et al.*, 2020; 2021), of which ~86% are described, and the remaining awaiting formal treatment are mostly 'known unknowns' (Ronco *et al.*, 2020).

In a test of the potential of DNA barcoding of the Tanganyikan assemblage, out of 96 species of littoral cichlids assayed, Breman *et al.* (2016) successfully *identified* only 73% of the valid species included. The potential for *discovery* was not much better, either underestimating by ~30% (70 hypothetical molecular operational taxonomic units (MOTUs) using the General Mixed Yule Coalescent) or overestimating species-level diversity by ~10% (115 hypothetical MOTUs using the Automatic Barcode Gap Discovery method). BINs are thus a poor proxy to species in the case of Tanganyikan cichlids.

The likely reasons for this poor behaviour of DNA barcoding in identifying or delineating Tanganyikan

cichlids reported by Breman et al. (2016) stem from a combination of introgression and incomplete lineage sorting (i.e., sharing of haplotypes across species and apparent polyphyly; e.g., Brawand et al., 2014; Gante et al., 2016; Nevado et al., 2011), associated with fast cladogenesis observed in speciose groups (i.e., lack of universal resolution of the barcode at the species level). The results certainly do not reflect major issues with species validity, as morphology and nuclear genomes both support the vast majority of the species recognized (Ronco et al., 2021). Thus, limitations of DNA barcoding are overcome with an integrative approach to identification and delineation.

Considering that these difficulties are readily apparent in the oldest of cichlid radiations, a minimalist DNA taxonomy would be deemed unfeasible in younger radiations such as those of lakes Malawi and Victoria, which are far more speciose and have been boosted by ancient hybridisation (Malinsky et al., 2018; Meier et al., 2017). Species boundaries permeable to introgression (in particular of mtDNA) and rapid cladogenesis are attributes of hyperdiverse groups. These are the groups for which Sharkey et al. (2021b) claim a minimalist DNA taxonomy would speed up naming of new species, although that would be done at the cost of rigour, accuracy, or reliability in identifying or delineating new species.

Butterflies, other insects, and the ‘Wolbachia effect’.

There is no other insect group so intensively studied as European butterflies, save perhaps *Drosophila* Fallén, 1823. Given the attractiveness of European butterflies and the long history of their study, we know a lot about their distributional changes, ecologies and life histories. This was followed by the emergence of national and international barcoding programs (e.g., Lukhtanov et al., 2009; Dincă et al., 2011; Hausmann et al., 2011; Dincă et al., 2015; Litman et al., 2018; Dincă et al., 2021). In this well-known group, species delineation methods revealed a portion of polyphyletic, paraphyletic or barcode-sharing species. For example, Dincă et al. (2011) showed that in 180 Romanian butterfly species, only 162 species formed separate barcode clusters, three species pairs shared barcodes, four species were paraphyletic, two species pairs were polyphyletic and eight species had intraspecific divergence (p-distances) over 2.0%, which, according to the minimalist approach, would merit their description as a complex of cryptic species for which morphology ‘fails’. Similar numbers were obtained for other countries (Hausmann et al., 2011; Dincă et al., 2015; Litman et al., 2018). Recently, a compilation of 22,306 sequences of 459 European butterfly species (97% of known European species) was analysed by Dincă et al. (2021). In this dataset, the 459

species were assigned to 441 BINs, in which 65 BINs were taxonomically discordant, and 16 were singleton (i.e., not including any other sequences), 69 species shared barcodes and 56 species had a divergence greater than 2.5%.

A comparable picture arose for the North American butterfly fauna (D’Ercole et al., 2021): in the 814 American butterfly species, 573 species (out of 755 with more than 2 specimens) formed distinct clusters, 151 species were polyphyletic or paraphyletic, 125 species shared barcodes and 79 species had a divergence greater than 2.5%. An analogous analysis of dragonflies, a genetically distant insect order, produced similar results (Geiger et al., 2021), indicating that such patterns of non-monophyly are widely distributed among insects, and they might be even much broader: in an often-cited review on mitochondrial phylogenies of closely related animal species groups, over 20% of ~2,300 species assayed showed species-level paraphyly or polyphyly (Funk & Omland, 2003). This would result in an unacceptably high percentage of incorrectly described species should only DNA barcodes be used to delineate species.

Mutanen et al. (2016) explored the amount of bias in European Lepidoptera in barcode paraphyly and polyphyly. They investigated 4,977 species (41,583 specimens) and found 465 non-monophyletic species using a neighbour-joining method and 469 species using the Maximum Likelihood method of tree reconstruction. These can be explained by over-splitting or cryptic speciation (estimated at 31.8%). However, they also recognized that non-monophyly increases when more specimens are investigated and is higher if the species are relatively young.

Such large-scale studies serve as a background for detection of cryptic diversity. The ideal procedure to follow them up with and produce new species identifications would include sequencing nuclear DNA, searching for morphological and life history traits, and (ideally) microbial screening. This array of techniques was used when delineating the European species of *Spialia* Swinhoe, 1912 (Hernández-Roldán et al., 2016). In many cases, the issues of barcode non-monophyly can be solved with detailed studies at intraspecific- and closely related species-group levels. Some butterfly species display mtDNA over-splitting: *Thymelicus sylvestris* (Poda, 1761) has six mitochondrial lineages, but these are not recovered by ddRADseq data (Hinojosa et al., 2019). An analogous pattern was observed in *Melitaea didyma* (Esper, 1778) (Dincă et al., 2019). *Melitaea ornata* Christoph, 1893 includes two separate mitochondrial groups and one of them clusters with its sister species *Melitaea phoebe* (Denis & Schiffermüller, 1775),

whereas nuclear genes clearly distinguish two species (Tóth *et al.*, 2017). Younger groups, where speciation is ongoing, can display incomplete lineage sorting, identifiable by an absence of a predictable biogeographical pattern (Funk & Omland, 2003; Toews & Brelsford, 2012). The introgression of mitochondrial DNA occurs via hybridisation, which is observable in the butterfly genus *Colias* Fabricius, 1807 (Wheat & Watt, 2008).

In addition, mitochondrial DNA inheritance patterns of arthropods and filarial nematodes are affected by the co-inherited bacteria *Wolbachia* Hertig, 1936 (e.g., Hurst & Jiggins, 2005). Ahmed *et al.* (2015) estimated the presence of endosymbiotic *Wolbachia* in 80% of Lepidoptera, with 25–33% infected individuals. As *Wolbachia* interferes with reproduction of its hosts, it might cause a swift spread through a population, hitchhiking one mitochondrial type. Mitochondrial clades with a deep divergence associated with the presence/absence of *Wolbachia* were observed in both *Phengaris nausithous* (Bergsträsser, 1779) and *P. teleius* (Bergsträsser, 1779) (Ritter *et al.*, 2013). If two bacterial strains are present, they could each bear a different mitotype, causing a deep mitochondrial split, as observed in American *Coenonympha tullia* (Müller, 1764) (Kodandaramaiah *et al.*, 2013). Furthermore, *Wolbachia* might mediate introgression, as hypothesised for *Iphioides podalirius* (Linnaeus, 1758) and *I. feisthamelii* (Duponchel, 1832) (Gaunet *et al.*, 2019). *Wolbachia* screening might support past hybridisation events as in *M. phoebe* and *M. ornata* (Tóth *et al.*, 2017), and *Pseudophilotes baton* (Bergsträsser, 1779) and *P. vicrama* (Moore, 1865) (Sucháčková Bartoňová *et al.*, 2021).

Large datasets, compiled and extended in Mutanen *et al.* (2016) and Dincă *et al.* (2021) demonstrate what would happen if the minimalist approach would have been used. The minimalist approach would have accepted barcode-delimited species largely as is, without taking into account morphological and life history traits (as in the examples above), studying intraspecific or closely related species group variation in detail, or checking for the effect that *Wolbachia* could have on results. In the majority of cases, barcodes succeed in assigning specimens into proper taxonomic species, but they still have their limits for a substantial portion of taxa. It is fair to assume that a similar portion of species would have been inaccurately delimited in other, less explored insect groups. We have to consider that the amount of non-monophyly reflects the number of investigated specimens and that it depends on species age, i.e., low divergence in young species and high divergence in old species (Mutanen *et al.*, 2016).

DNA barcode-based minimalist taxonomy does not address the taxonomic impediment

Are the false negative and false positive rates for cichlids, butterflies and dragonflies discussed above acceptable in the face of the pressing need to name vanishing biodiversity? One could argue that a large proportion would still be correctly named using only a DNA barcode. One problem with a minimalist DNA approach is that we would be completely blind as to which taxa might or might not be properly named. In other words, given the paucity of taxonomic information gathered by minimalist DNA taxonomy, invalid taxa would effectively be ‘unknown unknowns’ – we know they would exist but we would not easily know which ones they might be without an in-depth reanalysis of the entire group (see Meier *et al.*, 2022). Thus, the amount of taxonomic work needed to assess which species are actually valid would surmount the work needed for a proper first pass using an integrative approach, even considering incremental contributions and subsequent efforts as new taxa are discovered. Furthermore, it is doubtful that taxonomy end-uses, such as conservation and management actions, would gladly be based on such high numbers of erroneously named and improperly circumscribed species, in particular since we ought to preserve and manage morphological and ecological diversity as much as molecular (i.e., barcode) diversity.

A path forward

Much emphasis has been placed in automation of molecular analyses and DNA barcoding has been surfing this wave. Sequencing technologies have witnessed great progress, as we quickly moved from automated Sanger sequencing, to ‘next generation’ short read sequencing, to ‘third generation’ long read sequencing in just two decades. One recent application that combines barcoding and high throughput sequencing is metabarcoding of environmental DNA (eDNA) for biodiversity monitoring. Metabarcoding of eDNA allows the simultaneous assessment of barcodes of multiple taxa present in the environment, for example from water, soil or air samples, as evidence of organisms’ presence (Taberlet *et al.*, 2018). A newer metagenomics approach is being developed that extends this principle to any type of DNA present in the environment (i.e., beyond barcodes), with gains in detection sensitivity, accuracy and precision, even when applied to multicellular eukaryotes (e.g., Alves *et al.*, 2019; Curto *et al.*, in prep). These high throughput approaches enable not only detecting known biodiversity (i.e., formerly characterised barcodes or genomes in known, previously described species), they also show a great potential for *discovering* unknown

molecular diversity. Such environmental metagenomics data could sensibly direct follow-up integrative studies towards targeting the collection of potential new taxa (to which the newly found molecular diversity belongs to), effectively closing the gap between known and unknown biodiversity.

Much less focus has been put on one of the bottlenecks to taxonomic studies: morphological analyses. In particular, one area that could benefit from automation, speeding up informative species descriptions, and ameliorating the taxonomic impediment, would be automated morphological classification and character *discovery*. Over the last decades we have seen considerable advancements in deep learning computational methods such as convolutional neural networks (CNNs) and other types of networks (e.g., Gaston & O’Neil, 2004). In particular, CNNs learn to extract relevant features from photographs without human intervention (e.g., Kaya et al., 2019; Valan et al., 2019). On the one hand, these methods could speed up routine identifications of a large proportion of taxa, freeing taxonomists for more creative and critical tasks. Specimens that do not fit the groups being classified could be flagged for more in-depth identification by taxonomists, with the potential for discovering new groups and undescribed species (e.g., Yang et al., 2021). On the other hand, as CNNs are classifying individuals, they are ‘learning’ about traits that group similar individuals and separate them from other groups. Thus, these methods have a great potential for *discovering* traits that discriminate between different species, facilitating species descriptions and highlighting traits that an experienced and dedicated taxonomist can test for applicability, including of cryptic taxa for which traditional characters have low resolution; of course, the goal is not that CNNs replace an experienced taxonomist, but rather provide additional information (e.g. highlighting potentially useful traits, verbal descriptions of the traits) that would speed up the process and enrich taxonomic descriptions. An investment on rapid automated biodiversity assessment was also recently proposed by Ahrens et al. (2021).

Soon it might be possible to not have to make difficult choices between thoroughness and speed of taxonomic treatments, in particular if cost and technology continue progressing at a steady pace. For instance, robotic coupling of sorting and classifying, photographing, and sequencing barcodes (without destroying the specimens in question) has recently been implemented for very small insects (Srivathsan et al., 2021; Würl et al., 2021). The initial investment might be steep for individuals, but probably within the reach of institutions and research projects dedicated to monitoring and describing biodiversity. Progress in this area should

allow addressing the taxonomic impediment without sacrificing rigour, utility or quality for speed and quantity.

Bioinformatics could also play a major role in next-generation taxonomy and enhancement of species discovery. A promising example is a user-friendly, specimen-based toolkit called ‘iTaxoTools 0.1’ (Vences et al., 2021a). This software, based on open-source Python code, includes tools focusing on species delimitation and diagnosis and centred around specimen identifiers, and at its current stage contains GUI versions of six species delimitation programs (ABGD, ASAP, DELINEATE, GMYC, PTP, tr2) and a simple threshold-clustering delimitation tool, along with new Python implementations of existing algorithms, including tools to compute pairwise DNA distances, ultrametric time trees based on non-parametric rate smoothing, species-diagnostic nucleotide positions, standard morphometric analyses, etc. This innovation could greatly accelerate the rate of species discovery in hyperdiverse taxa in the future, without compromising the quality of the descriptions.

Another important aspect is the cost associated with taxonomic activities (either personnel, robotics for morphology, DNA barcoding, computing power, or collection management), in particular if we are to adopt new strategies to speed them up. In the current model, it is undeniable that funding goes where perceived publication impacts are, and these are increasingly being measured as paper citations, which in turn translate into journal impact factors. The most direct way of recognising the importance of taxonomic work would be to make citations of species descriptions or other relevant taxonomic treatments (as argued elsewhere by Meier, 2017) mandatory for each taxon the study focuses on. In a digital era, publication page limits and word limits should be extended to allow the citing of original descriptions or taxonomic revisions of focal species studied. Considering that taxonomic descriptions are scientific hypotheses (Pante et al., 2015), and that current scientific practice rightfully calls for citing all relevant literature (e.g., statistical software), it is only fair that taxonomic literature receives equal treatment. Increased funding is necessary to address shortages of trained taxonomists, and challenges faced by Natural History museums and the utilisation of the biological collections they hold (e.g., Ebach et al., 2011; Pinheiro et al., 2019; Dupérré, 2020; Engel et al., 2021).

A frequently hidden aspect of DNA barcoding reliability is that it is only as good as the assignment of names to barcoded units. That requires careful identification of the individuals that are barcoded, using original descriptions, subsequent revisions and keys, and

ideally also examination of original type material. That rigour is seldom practised or explicitly stated, especially in labs focussed mainly on genetic or genomic analysis, where morphological identification skills are not honed, and where more work is done with tissue samples than with actual specimens. Misidentification of the material that later winds up serving as the barcode reference for a species can cause knock-on errors for decades, until a mismatch between the type material and the application of the name is discovered. Even when a great deal of rigour has been applied to the original assignment of DNA barcodes to species names, DNA sequences from historical type material have resulted in surprises that have necessitated synonymisations and replacement names (e.g., Vences *et al.*, 2021b). In the absence of that rigour, such mistakes will certainly be more common and necessitate even more revisionary work. For this reason, we argue that vouchering of DNA barcoded specimens in public museum collections should be the norm.

Finally, as we have discussed above, morphological comparison with existing nomina can be difficult in truly cryptic clades, or where the historical type material is damaged, ambiguous, or immature. However, it is no longer the case that the assignment of such names must necessarily be left to convention or best guesses. Major strides have been made recently in sequencing historical type and non-type material, some well over a century old, or fixed in DNA-degrading chemicals like formalin, and often based on very small quantities of tissue, in what has been dubbed museomics (e.g. Erpenbeck *et al.*, 2016; Ruane & Austin, 2017; Evans *et al.*, 2019; McGuire *et al.* 2018; Rancilhac *et al.* 2020; Scherz *et al.*, 2020; Call *et al.*, 2021; Straube *et al.*, 2021a, 2021b; Vences *et al.*, 2021b). This means that DNA barcodes (and indeed substantial parts of the mitochondrial and nuclear genomes) can be sequenced from old types, and used to more confidently assign names to lineages. As discussed above (and argued by some of us elsewhere; Zamani *et al.*, 2021), assignment of previous names should take priority over the establishment of new names, in order to reduce the strain on the taxonomic community.

Conclusions

As already argued by Wheeler (2008), taxonomy (i.e., species description) is different from instant species identification and should be considered a science of its own, with its own set of hypotheses to test. However, minimalist taxonomy relegates taxonomy to a mere service for other sciences. How can we move forward from this? Do we really need to consider taxonomy just a

tool for the end-user, ignoring the epistemology that sets taxonomy apart from other sciences? Probably the best way forward is to set aside any baseless claim of superiority of one data type over another and focus instead on building communities with the intention to create a collaborative environment where researchers with different sets of skills and passions can thrive. However, this means setting aside ego and personal achievement for a greater good: the discovery and study of what is left of the Earth's biodiversity. Is this achievable? Our response to this is affirmative. The ichneumonid community, as an example, was formally reignited in 2019 in Basel (Klopfstein *et al.*, 2019) with the aim to reduce the taxonomic impediment of the group by gathering people with different skill sets. Time will tell if the community will last, but the simple fact that it has been built points to the fact that collaboration, sharing data, diversifying the fields of knowledge and building bridges are achievable and, even more importantly, desirable actions.

Acknowledgments

Comments and suggestions by Tapani E. Hopkins, David Williams (Associate Editor), Richard Bateman (Royal Botanic Gardens, Kew) and an anonymous reviewer are greatly appreciated.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the USDA National Institute of Food and Agriculture, 1890 Institution Capacity Building Grant Project 1021805 to A.B.O; and by Fundação para a Ciência e a Tecnologia (Lisbon) (F.C.T.) grant number PTDC/BIA-CBI/31644/2017 to H.F.G.

ORCID


Alireza Zamani  <http://orcid.org/0000-0002-8084-9666>

Davide Dal Pos  <http://orcid.org/0000-0002-9122-934X>

Zdenek Faltýnek Fric  <http://orcid.org/0000-0002-3611-8022>

Alexander B. Orfinger  <http://orcid.org/0000-0002-4907-3150>

Mark D. Scherz  <http://orcid.org/0000-0002-4613-7761>

Alena Sucháčková Bartoňová  <http://orcid.org/0000-0001-6298-2466>

Hugo F. Gante  <http://orcid.org/0000-0002-0321-3023>

References

- Abdala, C. S., Paz, M. M., Semhan, R. V., García, N., Aguilar-Kirigin, A. J., Fariás, M. E., Valladares, P., Poblete, R. G., Quipildor, M. A., Valdes, J., & Langstroth, R. (2021). Increasing knowledge of the denizens of saline environments through integrative taxonomy: New Argentinian endemic taxa of *Liolaemus* (Iguania: Liolaemidae) and their evolutionary relationships. *Systematics and Biodiversity*, *19*, 135–167. <https://doi.org/10.1080/14772000.2020.1844818>
- Ahmed, M. Z., Araujo-Jnr, E. V., Welch, J. J., & Kawahara, A. Y. (2015). *Wolbachia* in butterflies and moths: Geographic structure in infection frequency. *Frontiers in Zoology*, *12*, 16. <https://doi.org/10.1186/s12983-015-0107-z>
- Ahrens, D., Ah Yong, S. T., Ballerio, A., Barclay, M., Eberle, J., Espeland, M., Huber, B. A., Mengual, X., Pacheco, T. L., Peters, R., Rulik, B., Vaz-de-Mello, F., Wesener, T., & Krell, F.-T. (2021). Is it time to describe new species without diagnoses? A comment on Sharkey et al., (2021). *Zootaxa*, *5027*, 151–159.
- Alves, M. J., Gante, H., Veríssimo, A., Ribeiro, F., Santos, C. D., & Jentoft, S. (2019). From rare species detection to whole-community diversity using high-throughput sequencing of freshwater eDNA. *Biodiversity Information Science and Standards*, *3*, e37975. <https://doi.org/10.3897/biss.3.37975>
- Arthofer, W., Rauch, H., Thaler-Knoflach, B., Moder, K., Muster, C., Schlick-Steiner, B. C., & Steiner, F. M. (2013). How diverse is *Mitopus morio*? Integrative taxonomy detects cryptic species in a small-scale sample of a widespread harvestman. *Molecular Ecology*, *22*, 3850–3863. <https://doi.org/10.1111/mec.12340>
- Bond, J. E., Godwin, R. L., Colby, J. D., Newton, L. G., Zahnle, X. J., Agnarsson, I., Hamilton, C. A., & Kuntner, M. (2021). Improving Taxonomic Practices and Enhancing Its Extensibility—An Example from Araneology. *Diversity*, *14*, 5. <https://doi.org/10.3390/d14010005>
- Brasero, N., Martinet, B., Michez, D., Lecocq, T., Valterova, I., & Rasmont, P. (2020). Taxonomic revision of the *Sylvarum* group of bumblebees using an integrative approach. *Systematics and Biodiversity*, *18*, 12–28. <https://doi.org/10.1080/14772000.2020.1737843>
- Brawand, D., Wagner, C. E., Li, Y. I., Malinsky, M., Keller, I., Fan, S., Simakov, O., Ng, A. Y., Lim, Z. W., Bezault, E., Turner-Maier, J., Johnson, J., Alcazar, R., Noh, H. J., Russell, P., Aken, B., Alföldi, J., Amemiya, C., Azzouzi, N., ... Di Palma, F. (2014). The genomic substrate for adaptive radiation in African cichlid fish. *Nature*, *513*, 375–381. <https://doi.org/10.1038/nature13726>
- Breman, F. C., Loix, S., Jordaens, K., Snoeks, J., & Van Steenberge, M. (2016). Testing the potential of DNA barcoding in vertebrate radiations: The case of the littoral cichlids (Pisces, Perciformes, Cichlidae) from Lake Tanganyika. *Molecular Ecology Resources*, *16*, 1455–1464.
- Call, E., Mayer, C., Twort, V., Dietz, L., Wahlberg, N., & Espeland, M. (2021). Museomics: Phylogenomics of the moth family Epicopeiidae (Lepidoptera) using target enrichment. *Insect Systematics and Diversity*, *5*, 6. <https://doi.org/10.1093/isd/ixaa021>
- Carstens, B. C., Pelletier, T. A., Reid, N. M., & Satler, J. D. (2013). How to fail at species delimitation. *Molecular Ecology*, *22*, 4369–4383.
- Carvalho, M. R., Bockmann, F. A., Amorim, D. S., Brandão, C., Vivo, M., Figueiredo, J. L., Britski, H. A., Pinna, M., Menezes, N. A., Marques, F., Papavero, N., Cancellato, E. M., Crisci, J. V., McEachran, J. D., Schelly, R. S., Lundberg, J. G., Gill, A. G., Britz, R., Wheeler, Q. D., ... Nelson, G. J. (2007). Taxonomic impediment or impediment to taxonomy? A commentary on systematics and the cybertaxonomic-automation paradigm. *Evolutionary Biology*, *34*, 140–143. <https://doi.org/10.1007/s11692-007-9011-6>
- Chambers, E. A., & Hillis, D. M. (2020). The multispecies coalescent over-splits species in the case of geographically widespread taxa. *Systematic Biology*, *69*, 184–193. <https://doi.org/10.1093/sysbio/syz042>
- Chaplin, K., Sumner, J., Hipsley, C. A., & Melville, J. (2020). An integrative approach using phylogenomics and high-resolution x-ray computed tomography for species delimitation in cryptic taxa. *Systematic Biology*, *69*, 294–307. <https://doi.org/10.1093/sysbio/syz048>
- Costa, W., & Katz, A. M. (2021). Integrative taxonomy supports high species diversity of south-eastern Brazilian mountain catfishes of the *T. reinhardti* group (Siluriformes: Trichomycteridae). *Systematics and Biodiversity*, *19*, 601–621. <https://doi.org/10.1080/14772000.2021.1900947>
- Dal Pos, D., & Rouse, P. (2018). First record of the genus *Genaemirum* (Hymenoptera, Ichneumonidae, Ichneumoninae) from West Africa, with the description of a new species from Guinea. *European Journal of Taxonomy*, *473*, 1–11.
- Dayrat, B. (2005). Towards integrative taxonomy. *Biological Journal of the Linnean Society*, *85*, 407–415. <https://doi.org/10.1111/j.1095-8312.2005.00503.x>
- D’Ercole, J., Dincă, V., Opler, P. A., Kondla, N., Schmidt, C., Phillips, J. D., Robbins, R., Burns, J. M., Miller, S. E., Grishin, N., Zakharov, E. V., DeWaard, J. R., Ratnasingham, S., & Hebert, P. D. N. (2021). A DNA barcode library for the butterflies of North America. *PeerJ*, *9*, e11157.
- DeSalle, R., Egan, M. G., & Siddall, M. (2005). The unholy trinity: Taxonomy, species delimitation and DNA barcoding. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *360*, 1905–1916. <https://doi.org/10.1098/rstb.2005.1722>
- Di Giovanni, F., & Reshchikov, A. (2016). Contribution to the knowledge of Ichneumonidae (Hymenoptera) in Italy. *Linzer Biologische Beiträge*, *48*, 495–505.
- Di Giovanni, F., Reshchikov, A., Riedel, M., Diller, E., & Schwarz, M. (2015). New records of Ichneumonidae (Hymenoptera) for the Italian fauna. *Biodiversity Data Journal*, *3*, e5057. <https://doi.org/10.3897/BDJ.3.e5057>
- Di Giovanni, F., Scaramozzino, P. L., & Diller, E. (2018). A new species of *Misetus* Wesm., 1845 (Hymenoptera, Ichneumonidae, Ichneumoninae, Phaeogenini) from Italy, with updated key to the females of Palaearctic species. *Zootaxa*, *4374*, 594–600. <https://doi.org/10.11646/zootaxa.4374.4.8>

- Dincă, V., Dapporto, L., Somervuo, P., Vodă, R., Cuvelier, S., Gascoigne-Pees, M., Huemer, P., Mutanen, M., Hebert, P. D., & Vila, R. (2021). High resolution DNA barcode library for European butterflies reveals continental patterns of mitochondrial genetic diversity. *Communications Biology*, 4, 315. <https://doi.org/10.1038/s42003-021-01834-7>
- Dincă, V., Montagud, S., Talavera, G., Hernández-Roldán, J., Munguira, M. L., García-Barros, E., Hebert, P. D., & Vila, R. (2015). DNA barcode reference library for Iberian butterflies enables a continental-scale preview of potential cryptic diversity. *Scientific Reports*, 5, 12395. <https://doi.org/10.1038/srep12395>
- Dincă, V., Lee, K. M., Vila, R., & Mutanen, M. (2019). *The conundrum of species delimitation: A genomic perspective on a mitogenetically super-variable butterfly* [Paper presentation]. Proceedings of the Royal Society B: Biological Sciences, 286, 20191311.
- Dincă, V., Zakharov, E. V., Hebert, P. D., & Vila, R. (2011). *Complete DNA barcode reference library for a country's butterfly fauna reveals high performance for temperate Europe* [Paper presentation]. Proceedings of the Royal Society B: Biological Sciences, 278, 347–355.
- Downton, M., Meiklejohn, K., Cameron, S. L., & Wallman, J. (2014). A preliminary framework for DNA barcoding, incorporating the multispecies coalescent. *Systematic Biology*, 63, 639–644. <https://doi.org/10.1093/sysbio/syu028>
- Dupérré, N. (2020). Old and new challenges in taxonomy: What are taxonomists up against? *Megataxa*, 1, 59–62. <https://doi.org/10.11646/megataxa.1.1.12>
- Ebach, M. C., & Holdrege, C. (2005). DNA barcoding is no substitute for taxonomy. *Nature*, 434, 697. <https://doi.org/10.1038/434697b>
- Ebach, M. C., Valdecasas, A. G., & Wheeler, Q. D. (2011). Impediments to taxonomy and users of taxonomy: Accessibility and impact evaluation. *Cladistics: The International Journal of the Willi Hennig Society*, 27, 550–557. <https://doi.org/10.1111/j.1096-0031.2011.00348.x>
- Edwards, D. L., & Knowles, L. L. (2014). Species detection and individual assignment in species delimitation: Can integrative data increase efficacy? *Proceedings. Biological Sciences*, 281, 20132765. <https://doi.org/10.1098/rspb.2013.2765>
- Engel, M. S., Ceríaco, L. M. P., Daniel, G. M., Dellapé, P. M., Löbl, I., Marinov, M., Reis, R. E., Young, M. T., Dubois, A., Agarwal, I., Lehmann, A. P., Alvarado, M., Alvarez, N., Andreone, F., Araujo-Vieira, K., Ascher, J. S., Baêta, D., Baldo, D., Bandeira, S. A., ... Zacharie, C. K. (2021). The taxonomic impediment: A shortage of taxonomists, not the lack of technical approaches. *Zoological Journal of the Linnean Society*, 193, 381–387. <https://doi.org/10.1093/zoolinnean/zlab072>
- Erpenbeck, D., Ekins, M., Enghuber, N., Hooper, J., Lehnert, H., Polisenio, A., Schuster, A., Setiawan, E., De Voogd, N. J., Wörheide, G., & Van Soest, R. (2016). Nothing in (sponge) biology makes sense – except when based on holotypes. *Journal of the Marine Biological Association of the United Kingdom*, 96, 305–311. <https://doi.org/10.1017/S0025315415000521>
- Evans, B. J., Gansauge, M.-T., Stanley, E. L., Furman, B., Cauret, C., Ofori-Boateng, C., Gvoždík, V., Streicher, J. W., Greenbaum, E., Tinsley, R. C., Meyer, M., & Blackburn, D. C. (2019). *Xenopus fraseri*: Mr. Fraser, where did your frog come from? *PLoS One*, 14, 14, e0220892.
- Feliciano, K., Malaquias, M., Stout, C., Brenzinger, B., Gosliner, T. M., & Valdés, A. (2021). Molecular and morphological analyses reveal pseudocryptic diversity in *Micromelo undatus* (Bruguière, 1792) (Gastropoda: Heterobranchia: Aplustridae). *Systematics and Biodiversity*, 19, 834–858. <https://doi.org/10.1080/14772000.2021.1939458>
- Fernandez-Triana, J. L. (2022). Turbo taxonomy approaches: Lessons from the past and recommendations for the future based on the experience with Braconidae (Hymenoptera) parasitoid wasps. *ZooKeys*, 1087, 199–220. <https://doi.org/10.3897/zookeys.1087.76720>
- Funk, D. J., & Omland, K. E. (2003). Species-level paraphyly and polyphyly: Frequency, causes, and consequences, with insights from animal mitochondrial DNA. *Annual Review of Ecology, Evolution, and Systematics*, 34, 397–423. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132421>
- Gante, H. F., Matschiner, M., Malmström, M., Jakobsen, K. S., Jentoft, S., & Salzburger, W. (2016). Genomics of speciation and introgression in Princess cichlid fishes from Lake Tanganyika. *Molecular Ecology*, 25, 6143–6161.
- Gante, H. F., & Salzburger, W. (2012). Evolution: Cichlid models on the runaway to speciation. *Current Biology: CB*, 22, R956–R958.
- Gaston, K. J., & O'Neill, M. A. (2004). Automated species identification: Why not? *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359, 655–667. <https://doi.org/10.1098/rstb.2003.1442>
- Gaunet, A., Dincă, V., Dapporto, L., Montagud, S., Vodă, R., Schär, S., Badiane, A., Font, E., & Vila, R. (2019). Two consecutive *Wolbachia*-mediated mitochondrial introgressions obscure taxonomy in Palearctic swallowtail butterflies (Lepidoptera, Papilionidae). *Zoologica Scripta*, 48, 507–519. <https://doi.org/10.1111/zsc.12355>
- Gebiola, M., Gómez-Zurita, J., Monti, M. M., Navone, P., & Bernardo, U. (2012). Integration of molecular, ecological, morphological and endosymbiont data for species delimitation within the *Pnigalio soemius* complex (Hymenoptera: Eulophidae). *Molecular Ecology*, 21, 1190–1208. <https://doi.org/10.1111/j.1365-294X.2011.05428.x>
- Geiger, M., Koblmüller, S., Assandri, G., Chovanec, A., Ekrem, T., Fischer, I., Galimberti, A., Grabowski, M., Haring, E., Hausmann, A., Hendrich, L., Koch, S., Mamos, T., Rothe, U., Rulik, B., Rewicz, T., Sittenthaler, M., Stur, E., Tończyk, G., Zangl, L., & Moriniere, J. (2021). Coverage and quality of DNA barcode references for Central and Northern European Odonata. *PeerJ*, 9, e11192.
- Gregory, T. R. (2005). DNA barcoding does not compete with taxonomy. *Nature*, 434, 1067. <https://doi.org/10.1038/4341067b>
- Hausmann, A., Haszprunar, G., Segerer, A. H., Speidel, W., Behounek, G., & Hebert, P. D. (2011). Now DNA-barcode: The butterflies and larger moths of Germany. *Spixiana*, 34, 47–58.
- Hebert, P., & Barrett, R. (2005). Reply to the comment by L. Prendini on “Identifying spiders through DNA barcodes. *Canadian Journal of Zoology*, 83, 505–506. <https://doi.org/10.1139/z05-026>
- Hebert, P., & Gregory, T. R. (2005). The promise of DNA barcoding for taxonomy. *Systematic Biology*, 54, 852–859.
- Hebert, P., Cywinska, A., Ball, S. L., & Dewaard, J. R. (2003). *Biological identifications through DNA barcodes* [Paper presentation]. Proceedings of the Royal Society B: Biological Sciences, 270, 313–321.

- Heinrich, G. (1967). *Synopsis and reclassification of the Ichneumoninae Stenopneusticae of Africa south of Sahara (Hym.)* (Vol. 1, pp. 250). Farmington State College Press.
- Hernández-Roldán, J. L., Dapporto, L., Dincă, V., Vicente, J. C., Hornett, E. A., Šichová, J., Lukhtanov, V. A., Talavera, G., & Vila, R. (2016). Integrative analyses unveil speciation linked to host plant shift in *Spialia* butterflies. *Molecular Ecology*, 25, 4267–4284. <https://doi.org/10.1111/mec.13756>
- Hinojosa, J. C., Koubínová, D., Szenteczki, M. A., Pitteloud, C., Dincă, V., Alvarez, N., & Vila, R. (2019). A mirage of cryptic species: Genomics uncover striking mitonuclear discordance in the butterfly *Thymelicus sylvestris*. *Molecular Ecology*, 28, 3857–3868.
- Hopkins, T., Roininen, H., van Noort, S., Broad, G. R., Kaunisto, K., & Sääksjärvi, I. E. (2019). Extensive sampling and thorough taxonomic assessment of Afrotropical Rhyssinae (Hymenoptera, Ichneumonidae) reveals two new species and demonstrates the limitations of previous sampling efforts. *ZooKeys*, 878, 33–71. <https://doi.org/10.3897/zookeys.878.37845>
- Hurst, G., & Jiggins, F. M. (2005). *Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: The effects of inherited symbionts* [Paper presentation]. Proceedings of the Royal Society B: Biological Sciences, 272, 1525–1534.
- ICZN (1999). *International Code of Zoological Nomenclature* (4th ed). International Trust for Zoological Nomenclature.
- Johansson, N. (2020). Additions to the Swedish fauna of Diplazontinae (Hymenoptera: Ichneumonidae) with the descriptions of five new species. *European Journal of Taxonomy*, 724, 70–92. <https://doi.org/10.5852/ejt.2020.724.1159>
- Kaya, A., Keceli, A. S., Catal, C., Yalic, H. Y., Temucin, H., & Tekinerdogan, B. (2019). Analysis of transfer learning for deep neural network based plant classification models. *Computers and Electronics in Agriculture*, 158, 20–29. <https://doi.org/10.1016/j.compag.2019.01.041>
- Klopfstein, S. (2014). Revision of the western palaeartic diplazontinae (Hymenoptera, Ichneumonidae). *Zootaxa*, 3801, 1–143. <https://doi.org/10.11646/zootaxa.3801.1.1>
- Klopfstein, S., Santos, B. F., Shaw, M. R., Alvarado, M., Bennett, A., Dal Pos, D., Giannotta, M., Herrera, A., Karlsson, D., Khalaim, A. I., Lima, A. R., Mikó, I., Sääksjärvi, I. E., Shimizu, S., Spasojevic, T., van Noort, S., Vilhelmsen, L., & Broad, G. R. (2019). Darwin wasps: A new name heralds renewed efforts to unravel the evolutionary history of Ichneumonidae. *Entomological Communications*, 1, ec010. <https://doi.org/10.37486/2675-1305.ec01006>
- Kodandaramaiah, U., Simonsen, T. J., Bromilow, S., Wahlberg, N., & Sperling, F. (2013). Deceptive single-locus taxonomy and phylogeography: *Wolbachia*-associated divergence in mitochondrial DNA is not reflected in morphology and nuclear markers in a butterfly species. *Ecology and Evolution*, 3, 5167–5176. <https://doi.org/10.1002/ece3.886>
- Lima, L. R., Bruschi, D. P., Nascimento, F., Araújo, P., Costa, L. P., Thomé, M., Garda, A. A., Zattera, M. L., & Mott, T. (2020). Below the waterline: Cryptic diversity of aquatic pipid frogs (*Pipa carvalhoi*) unveiled through an integrative taxonomy approach. *Systematics and Biodiversity*, 18, 771–783. <https://doi.org/10.1080/14772000.2020.1795742>
- Lipscomb, D., Platnick, N., & Wheeler, Q. (2003). The intellectual content of taxonomy: A comment on DNA taxonomy. *Trends in Ecology & Evolution*, 18, 65–66. [https://doi.org/10.1016/S0169-5347\(02\)00060-5](https://doi.org/10.1016/S0169-5347(02)00060-5)
- Litman, J., Chittaro, Y., Birrer, S., Praz, C., Wermeille, E., Fluri, M., Stalling, T., Schmid, S., Wyler, S., & Gonthier, Y. (2018). A DNA barcode reference library for Swiss butterflies and forest moths as a tool for species identification, systematics and conservation. *PLoS One*, 13, e0208639. <https://doi.org/10.1371/journal.pone.0208639>
- Lukhtanov, V. A., Sourakov, A., Zakharov, E. V., & Hebert, P. D. (2009). DNA barcoding Central Asian butterflies: Increasing geographical dimension does not significantly reduce the success of species identification. *Molecular Ecology Resources*, 9, 1302–1310.
- Malinsky, M., Svardal, H., Tyers, A. M., Miska, E. A., Genner, M. J., Turner, G. F., & Durbin, R. (2018). Whole-genome sequences of Malawi cichlids reveal multiple radiations interconnected by gene flow. *Nature Ecology & Evolution*, 2, 1940–1955. <https://doi.org/10.1038/s41559-018-0717-x>
- Mason, N. A., Fletcher, N. K., Gill, B. A., Funk, W. C., & Zamudio, K. R. (2020). Coalescent-based species delimitation is sensitive to geographic sampling and isolation by distance. *Systematics and Biodiversity*, 18, 269–280. <https://doi.org/10.1080/14772000.2020.1730475>
- McGuire, J. A., Cotoras, D. D., O'Connell, B., Lawalata, S. Z. S., Wang-Claypool, C. Y., Stubbs, A., Huang, X., Wogan, G. O. U., Hykin, S. M., Reilly, S. B., Bi, K., Riyanto, A., Arida, E., Smith, L. L., Milne, H., Streicher, J. W., & Iskandar, D. T. (2018). Squeezing water from a stone: High-throughput sequencing from a 145-year old holotype resolves (barely) a cryptic species problem in flying lizards. *PeerJ*, 6, e4470. <https://doi.org/10.7717/peerj.4470>
- Meier, R. (2017). Citation of taxonomic publications: The why, when, what and what not. *Systematic Entomology*, 42, 301–304. <https://doi.org/10.1111/syen.12215>
- Meier, R., Blaimer, B. B., Buenaventura, E., Hartop, E., von Rintelen, T., Srivathsan, A., & Yeo, D. (2022). A re-analysis of the data in Sharkey et al.'s (2021) minimal list revision reveals that BINs do not deserve names, but BOLD Systems needs a stronger commitment to open science. *Cladistics: The International Journal of the Willi Hennig Society*, 38, 264–275. <https://doi.org/10.1111/cla.12489>
- Meier, J. I., Marques, D., Mwaiko, S., Wagner, C. E., Excoffier, L., & Seehausen, O. (2017). Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nature Communications*, 8, 14363. <https://doi.org/10.1038/ncomms14363>
- Meierotto, S., Sharkey, M. J., Janzen, D. H., Hallwachs, W., Hebert, P. D., Chapman, E. G., & Smith, M. A. (2019). A revolutionary protocol to describe understudied hyperdiverse taxa and overcome the taxonomic impediment. *Deutsche Entomologische Zeitschrift*, 66, 119–145. <https://doi.org/10.3897/dez.66.34683>
- Mora, C., Tittensor, D. P., Adl, S., Simpson, A., & Worm, B. (2011). How many species are there on earth and in the ocean? *PLoS Biology*, 9, e1001127.
- Moritz, C., & Cicero, C. (2004). DNA barcoding: Promise and pitfalls. *PLoS Biology*, 2, e279. <https://doi.org/10.1371/journal.pbio.0020354>
- Mutanen, M., Kivelä, S. M., Vos, R. A., Doorenweerd, C., Ratnasingham, S., Hausmann, A., Huemer, P., Dincă, V., van Nieukerken, E. J., Lopez-Vaamonde, C., Vila, R., Aarvik, L., Decaëns, T., Efetov, K. A., Hebert, P. D.,

- Johnsen, A., Karsholt, O., Pentinsaari, M., Rougerie, R., ... Godfray, H. C. (2016). Species-level para- and polyphyly in DNA barcode gene trees: Strong operational bias in European Lepidoptera. *Systematic Biology*, *65*, 1024–1040. <https://doi.org/10.1093/sysbio/syw044>
- Nevado, B., Fazalova, V., Backeljau, T., Hanssens, M., & Verheyen, E. (2011). Repeated unidirectional introgression of nuclear and mitochondrial DNA between four congeneric Tanganyikan cichlids. *Molecular Biology and Evolution*, *28*, 2253–2267. <https://doi.org/10.1093/molbev/msr043>
- Newton, L. G., Starrett, J., Hendrixson, B. E., Derkarabetian, S., & Bond, J. E. (2020). Integrative species delimitation reveals cryptic diversity in the southern Appalachian *Antrodiaetus unicolor* (Araneae: Antrodiaetidae) species complex. *Molecular Ecology*, *29*, 2269–2287. <https://doi.org/10.1111/mec.15483>
- Novacek, M. J., & Wheeler, Q. D. (1992). Extinct taxa—accounting for 99.9990...% of the earth's biota. In M. J. Novacek, & Q. D. Wheeler, (Eds.), *Extinction and phylogeny* (pp. 1–16). Columbia University Press.
- Olave, M., Solà, E., & Knowles, L. L. (2014). Upstream analyses create problems with DNA-based species delimitation. *Systematic Biology*, *63*, 263–271. <https://doi.org/10.1093/sysbio/syt106>
- Pante, E., Puillandre, N., Viricel, A., Arnaud, Haond, S., Aurelle, D., Castelin, M., Chenuil, A., Destombe, C., Forcioli, D., Valero, M., Viard, F., & Samadi, S. (2015). Species are hypotheses: Avoid connectivity assessments based on pillars of sand. *Molecular Ecology*, *24*, 525–544. <https://doi.org/10.1111/mec.13048>
- Papakostas, S., Michaloudi, E., Proios, K., Brehm, M., Verhage, L., Rota, J., Peña, C., Stamou, G., Pritchard, V. L., Fontaneto, D., & Declerck, S. (2016). Integrative taxonomy recognizes evolutionary units despite widespread mitonuclear discordance: Evidence from a rotifer cryptic species complex. *Systematic Biology*, *65*, 508–524. <https://doi.org/10.1093/sysbio/syw016>
- Pham, N. T., Matsumoto, R., Konishi, K., Sheng, M.-L., & Broad, G. R. (2020). A review of the genus *Apophysius* Cushman (Ichneumonidae: Phygadeuontinae), with descriptions of six new species. *Zootaxa*, *4802*, 301–316. <https://doi.org/10.11646/zootaxa.4802.2.5>
- Pinheiro, H. T., Moreau, C. S., Daly, M., & Rocha, L. A. (2019). Will DNA barcoding meet taxonomic needs? *Science (New York, N.Y.)*, *365*, 873–874. <https://doi.org/10.1126/science.aay7174>
- Puillandre, N., Modica, M. V., Zhang, Y., Sirovich, L., Boisselier, M.-C., Cruaud, C., Holford, M., & Samadi, S. (2012). Large-scale species delimitation method for hyperdiverse groups. *Molecular Ecology*, *21*, 2671–2691. <https://doi.org/10.1111/j.1365-294X.2012.05559.x>
- Rancilhac, L., Bruy, T., Scherz, M. D., Pereira, E. A., Preick, M., Straube, N., Lyra, M., Ohler, A., Streicher, J. W., Andreone, F., Crottini, A., Hutter, C. R., Randrianantoandro, J. C., Rakotoarison, A., Glaw, F., Hofreiter, M., & Vences, M. (2020). Target-enriched DNA sequencing from historical type material enables a partial revision of the Madagascar giant stream frogs (genus *Mantidactylus*). *Journal of Natural History*, *54*, 87–118. <https://doi.org/10.1080/00222933.2020.1748243>
- Ratnasingham, S., & Hebert, P. D. (2007). BOLD: The barcode of life data system (<http://www.barcodinglife.org>). *Molecular Ecology Notes*, *7*, 355–364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Ritter, S., Michalski, S. G., Settele, J., Wiemers, M., Fric, Z. F., Sielezniew, M., Sasic, M., Rozier, Y., & Durka, W. (2013). *Wolbachia* infections mimic cryptic speciation in two parasitic butterfly species, *Phengaris teleius* and *P. nausithous* (Lepidoptera: Lycaenidae). *PLoS One*, *8*, e78107. <https://doi.org/10.1371/journal.pone.0078107>
- Ronco, F., Büscher, H. H., Indermaur, A., & Salzburger, W. (2020). The taxonomic diversity of the cichlid fish fauna of ancient Lake Tanganyika, East Africa. *Journal of Great Lakes Research*, *46*, 1067–1078. <https://doi.org/10.1016/j.jglr.2019.05.009>
- Ronco, F., Matschiner, M., Böhne, A., Boila, A., Büscher, H. H., El Taher, A., Indermaur, A., Malinsky, M., Ricci, V., Kahmen, A., Jentoft, S., & Salzburger, W. (2021). Drivers and dynamics of a massive adaptive radiation in cichlid fishes. *Nature*, *589*, 76–81. <https://doi.org/10.1038/s41586-020-2930-4>
- Rousse, P., Broad, G. R., & van Noort, S. (2016). Review of the genus *Genaemirum* Heinrich (Hymenoptera, Ichneumonidae, Ichneumoninae) with interactive identification keys to species. *ZooKeys*, *636*, 77–105.
- Rousse, P., & van Noort, S. (2014). A review of the Afrotropical Rhyssinae (Hymenoptera: Ichneumonidae) with the descriptions of five new species. *European Journal of Taxonomy*, *91*, 1–42.
- Rousse, P., van Noort, S., & Diller, E. (2013). Revision of the Afrotropical Phaeogenini (Ichneumonidae, Ichneumoninae), with description of a new genus and twelve new species. *ZooKeys*, *354*, 1–85. <https://doi.org/10.3897/zookeys.354.5968>
- Ruane, S., & Austin, C. C. (2017). Phylogenomics using formalin-fixed and 100+ year-old intractable natural history specimens. *Molecular Ecology Resources*, *17*, 1003–1008. <https://doi.org/10.1111/1755-0998.12655>
- Scherz, M. D., Rasolonjatovo, S. M., Köhler, J., Rancilhac, L., Rakotoarison, A., Raselimanana, A. P., Ohler, A., Preick, M., Hofreiter, M., Glaw, F., & Vences, M. (2020). 'Barcode fishing' for archival DNA from historical type material overcomes taxonomic hurdles, enabling the description of a new frog species. *Scientific Reports*, *10*, 19109. <https://doi.org/10.1038/s41598-020-75431-9>
- Seberg, O., Humphries, C. J., Knapp, S., Stevenson, D. W., Petersen, G., Scharff, N., & Andersen, M. (2003). Shortcuts in systematics? A commentary on DNA-based taxonomy. *Trends in Ecology & Evolution*, *18*, 63–65. [https://doi.org/10.1016/S0169-5347\(02\)00059-9](https://doi.org/10.1016/S0169-5347(02)00059-9)
- Selfa, J., & Diller, E. (1994). Illustrated Key to the Western Palearctic Genera of Phaeogenini (Hymenoptera, Ichneumonidae, Ichneumoninae). *Entomofauna*, *15*, 237–252.
- Sharkey, M. J., Brown, B., Baker, A., & Mutanen, M. (2021a). Response to Zamani et al. (2020): The omission of critical data in the pursuit of “revolutionary” methods to accelerate the description of species. *ZooKeys*, *1033*, 191–201. <https://doi.org/10.3897/zookeys.1033.66186>
- Sharkey, M. J., Janzen, D. H., Hallwachs, W., Chapman, E. G., Smith, M. A., Dapkey, T., Brown, A., Ratnasingham, S., Naik, S., Manjunath, R., Perez, K., Milton, M., Hebert, P., Shaw, S. R., Kittel, R. N., Solis, M. A., Metz, M. A., Goldstein, P. Z., Brown, J. W., ... Burns, J. M. (2021b). Minimalist revision and description of 403 new species in 11 subfamilies of Costa Rican braconid parasitoid wasps, including host records for 219 species. *ZooKeys*, *1013*, 1–665. <https://doi.org/10.3897/zookeys.1013.55600>

- Srivathsan, A., Lee, L., Katoh, K., Hartop, E., Kutty, S. N., Wong, J., Yeo, D., & Meier, R. (2021). ONTbarcoder and MinION barcodes aid biodiversity discovery and identification by everyone, for everyone. *BMC Biology*, *19*, 217. <https://doi.org/10.1186/s12915-021-01141-x>
- Stork, N. E. (1997). *Measuring global biodiversity and its decline*. In M. L. Reaka-Kudla, D. E. Wilson, & E. O. Wilson (Eds.), *Biodiversity II: Understanding and protecting our biological resources* (pp. 41–48). Joseph Henry Press.
- Straube, N., Lyra, M. L., Paijmans, J., Preick, M., Basler, N., Penner, J., Rödel, M.-O., Westbury, M. V., Haddad, C. F., Barlow, A., & Hofreiter, M. (2021a). Successful application of ancient DNA extraction and library construction protocols to museum wet collection specimens. *Molecular Ecology Resources*, *21*, 2299–2315. <https://doi.org/10.1111/1755-0998.13433>
- Straube, N., Preick, M., Naylor, G., & Hofreiter, M. (2021b). Mitochondrial DNA sequencing of a wet-collection syntype demonstrates the importance of type material as genetic resource for lantern shark taxonomy (Chondrichthyes: Etmopteridae). *Royal Society Open Science*, *8*, 210474. <https://doi.org/10.1098/rsos.210474>
- Sucháčková Bartoňová, A., Konvička, M., Marešová, J., Wiemers, M., Ignatev, N., Wahlberg, N., Schmitt, T., & Fric, Z. F. (2021). *Wolbachia* affects mitochondrial population structure in two systems of closely related Palaearctic blue butterflies. *Scientific Reports*, *11*, 3019. <https://doi.org/10.1038/s41598-021-82433-8>
- Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For biodiversity research and monitoring*. Oxford University Press.
- Tereshkin, A. M. (2009). Illustrated key to the tribes of subfamilia Ichneumoninae and genera of the tribe Platylabini of world fauna (Hymenoptera, Ichneumonidae). *Linzer Biologische Beiträge*, *41*, 1317–1608.
- Toews, D., & Brelsford, A. (2012). The biogeography of mitochondrial and nuclear discordance in animals. *Molecular Ecology*, *21*, 3907–3930.
- Tóth, J. P., Varga, Z., Verovnik, R., Wahlberg, N., Váradi, A., & Bereczki, J. (2017). Mito-nuclear discordance helps to reveal the phylogeographic patterns of *Melitaea ornata* (Lepidoptera: Nymphalidae). *Biological Journal of the Linnean Society*, *121*, 267–281. <https://doi.org/10.1093/biolinnean/blw037>
- Valan, M., Makonyi, K., Maki, A., Vondráček, D., & Ronquist, F. (2019). Automated taxonomic identification of insects with expert-level accuracy using effective feature transfer from convolutional networks. *Systematic Biology*, *68*, 876–895. <https://doi.org/10.1093/sysbio/syz014>
- Varga, O. (2020). First record of *Epirhyssa* Cresson, 1865 (Hymenoptera, Ichneumonidae, Rhyssinae) from Kenya. *Zootaxa*, *4852*, zootaxa.4852.5.9–593. <https://doi.org/10.11646/zootaxa.4852.5.9>
- Vences, M., Köhler, J., Andreone, F., Craul, A.-K., Crottini, A., du Preez, L., Preick, M., Rancilhac, L., Rödel, M.-O., Scherz, M. D., Streicher, J. W., Hofreiter, M., & Glaw, F. (2021b). Target-enriched DNA sequencing clarifies the identity of name-bearing types of the *Gephyromantis plicifer* complex and reveals a new species of mantellid frog from Madagascar (Amphibia, Anura). *Spixiana*, *44*, 175–202.
- Vences, M., Miralles, A., Brouillet, S., Ducasse, J., Fedosov, A., Kharchev, V., Kostadinov, I., Kumari, S., Patmanidis, S., Scherz, M. D., Puillandre, N., & Renner, S. S. (2021a). iTaxoTools 0.1: Kickstarting a specimen-based software toolkit for taxonomists. *Megataxa*, *6*, 77–92. <https://doi.org/10.11646/megataxa.6.2.1>
- Vieites, D. R., Wollenberg, K. C., Andreone, F., Köhler, J., Glaw, F., & Vences, M. (2009). *Vast underestimation of Madagascar's biodiversity evidenced by an integrative amphibian inventory* [Paper presentation]. Proceedings of the National Academy of Sciences of the United States of America, 106, 8267–8272.
- Wachter, G. A., Muster, C., Arthofer, W., Rasputnig, G., Föttinger, P., Komposch, C., Steiner, F. M., & Schlick-Steiner, B. C. (2015). Taking the discovery approach in integrative taxonomy: Decrypting a complex of narrow-endemic Alpine harvestmen (Opiliones: Phalangidae: *Megabunus*). *Molecular Ecology*, *24*, 863–889. <https://doi.org/10.1111/mec.13077>
- Weston, J., Peart, R. A., & Jamieson, A. J. (2020). Amphipods from the Wallaby-Zenith Fracture Zone, Indian Ocean: New genus and two new species identified by integrative taxonomy. *Systematics and Biodiversity*, *18*, 57–78. <https://doi.org/10.1080/14772000.2020.1729891>
- Wheat, C. W., & Watt, W. B. (2008). A mitochondrial-DNA-based phylogeny for some evolutionary-genetic model species of *Colias* butterflies (Lepidoptera, Pieridae). *Molecular Phylogenetics and Evolution*, *47*, 893–902. <https://doi.org/10.1016/j.ympev.2008.03.013>
- Wheeler, Q. (2018). Blank Canvas: The case for descriptive taxonomy. *Integrative and Comparative Biology*, *58*, 1118–1121. <https://doi.org/10.1093/icb/icy067>
- Wheeler, Q. D. (2008). Undisciplined thinking: Morphology and Hennig's unfinished revolution. *Systematic Entomology*, *33*, 2–7. <https://doi.org/10.1111/j.1365-3113.2007.00411.x>
- Wheeler, Q. D. (2009). Revolutionary thoughts on taxonomy: Declarations of independence and interdependence. *Zoologia (Curitiba)*, *26*, 1–4. <https://doi.org/10.1590/S1984-46702009000100001>
- Wheeler, Q. D., Raven, P. H., & Wilson, E. O. (2004). Taxonomy: Impediment or expedient? *Science (New York, N.Y.)*, *303*, 285. <https://doi.org/10.1126/science.303.5656.285>
- Wiemers, M., & Fiedler, K. (2007). Does the DNA barcoding gap exist? A case study in blue butterflies (Lepidoptera: Lycaenidae). *Frontiers in Zoology*, *4*, 8–16. <https://doi.org/10.1186/1742-9994-4-8>
- Wiens, J. J. (2004). The role of morphological data in phylogeny reconstruction. *Systematic Biology*, *53*, 653–661. <https://doi.org/10.1080/10635150490472959>
- Will, K. W., Mishler, B. D., & Wheeler, Q. D. (2005). The perils of DNA barcoding and the need for integrative taxonomy. *Systematic Biology*, *54*, 844–851. <https://doi.org/10.1080/10635150500354878>
- Wüthrich, L., Pylatiuk, C., Giersch, M., Lapp, F., von Rintelen, T., Balke, M., Schmidt, S., Cerretti, P., & Meier, R. (2021). DiversityScanner: Robotic discovery of small invertebrates with machine learning methods. *BioRxiv*.
- Yang, B., Zhang, Z., Yang, C., Wang, Y., Orr, M. C., Hongbin, W., & Zhang, A.-B. (2021). Identification of species by combining molecular and morphological data using convolutional neural networks. *Systematic Biology*, <https://doi.org/10.1093/sysbio/syab076>
- Zamani, A., Fric, Z. F., Gante, H. F., Hopkins, T., Orfinger, A. B., Scherz, M. D., Sucháčková Bartoňová, A., & Pos,

- D. D. (2022). DNA barcodes on their own are not enough to describe a species. *Systematic Entomology*. <https://doi.org/10.1111/syen.12538>
- Zamani, A., Vahtera, V., Sääksjärvi, I. E., & Scherz, M. D. (2021). The omission of critical data in the pursuit of 'revolutionary' methods to accelerate the description of species. *Systematic Entomology*, 46, 1–4. <https://doi.org/10.1111/syen.12444>

Associate Editor: Dr. David Williams